



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

cloudy weather during January and February made it impossible to follow this object as closely as was desirable. Observations were secured with the 12-inch on January 13th and 15th, and with the 36-inch on February 15th. The observed places are in good agreement with those computed from the elliptic elements.

On February 15th the comet was a very difficult object to measure, even with the great refractor, and on March 8th it could not be seen at all, though the conditions were good.

Comet Brorsen, which was expected to return to perihelion in January, was looked for with the 36-inch telescope on several mornings in February, but without success. The atmospheric conditions during the search were good, and a field about two degrees square was examined on each night.

R. G. AITKEN.

SOME STARS WITH LARGE RADIAL VELOCITIES.

While pursuing the regular programme of observation with the Mills spectrograph, it was found that the following stars have large velocities in the line of sight, as indicated below:—

ϵ *Andromedæ* ($\alpha = 0^h 33^m$; $\delta = + 28^\circ 46'$).

1898, October 4	— 83.4 ^{km}	WRIGHT.
October 9	— 83.3	WRIGHT.
1899, August 29	— 84.6	WRIGHT.
1900, August 22	— 83.4	WRIGHT.
Mean,	— 83.7	

μ *Cassiopeiæ* ($\alpha = 1^h 0^m$; $\delta = + 54^\circ 20'$).

1900, September 9	— 97.2 ^{km}	WRIGHT.
September 18	— 97.0	WRIGHT.
December 11	— 98.	CAMPBELL.

The proper motion of μ *Cassiopeiæ* is 3".75 per year. JACOBY'S parallax, determined from the Rutherford photographs, is 0".275. These correspond to a motion at right angles to the line of sight of 66^{km} per second, though this includes nearly the full component of the motion of the solar system.

δ *Leporis* ($\alpha = 5^h 47^m.0$; $\delta = - 20^\circ 54'$).

1900, December 24	+ 95 ^{km}	CAMPBELL.
December 25	+ 96	CAMPBELL.
December 30	+ 94	CAMPBELL.

θ *Canis Majoris* ($\alpha = 6^h 50^m$; $\delta = -11^\circ 55'$).

1897, December	15	+ 96 ^{km}	CAMPBELL.
1899, October	16	+ 96.0	WRIGHT.
1900, October	9	+ 95.5	WRIGHT.

ι *Pegasi* ($\alpha = 21^h 17^m$; $\delta = +19^\circ 23'$).

1900, July	3	- 75.7 ^{km}	WRIGHT.
July	8	- 74.9	WRIGHT.
July	16	- 77.1	WRIGHT.

μ *Sagittarii* ($\alpha = 18^h 7^m.8$; $\delta = -21^\circ 05'$).

1899, June	19	- 75 ^{km}	WRIGHT.
1900, May	30	- 76	WRIGHT.

These measures are subject to an uncertainty of several kilometers, on account of the character of the spectrum.

The negative sign indicates approach toward the solar system, and the positive sign recession from the solar system.

W. W. CAMPBELL.

NOTE ON THE PROBABLE ERROR OF MICROMETER MEASURES OF *EROS*.

During the recent *Eros* campaign for the determination of the value of the solar parallax, a great many observations were made visually with the micrometer. The question of the accuracy of these measures is therefore an important one. In general, the motion of *Eros* was so rapid in both coordinates that the probable error of the mean of a number of settings, even when made in rapid succession, could not be determined in the usual manner by treating the differences between the single settings and the mean as the v 's.

But on December 5, 1900, *Eros* ceased its apparent westward motion among the stars and began to move eastward. Consequently on that date its motion in Right Ascension was very slight—only about twenty-two seconds of arc in eight hours.

On the night mentioned, I measured the position of *Eros* with respect to two small stars in the early evening, when it was from three to two hours east of the meridian, and with respect to two other stars later in the night, when it was from four to five hours west of the meridian. In all I made eighteen measures of difference of Right Ascension, each consisting of ten settings with the micrometer. As the ten settings in no case required